Feedback Control as a Framework for Understanding Tradeoffs in Biology

Noah J. Cowan[1,*], M. Mert Ankarali[1], Jonathan[2] P. Dyhr, Manu S. Madhav[1], Eatai Roth[2], Shahin Sefati[1], Simon Sponberg[2], Sarah A. Stamper[1], Eric S. Fortune[3] and Thomas L. Daniel[2]

[1]Department of Mechanical Engineering, Johns Hopkins University, Baltimore, MD 21218, USA, [2]Department of Biology, University of Washington, Seattle, WA 98195, USA, and [3]Department of Biological Sciences, New Jersey Institute of Technology, Newark, NJ 07102, USA
[*]Author for correspondence (ncowan@jhu.edu)

arXiv:1402.5702v1 [q-bio.QM] 24 Feb 2014

Abstract

Control theory arose from a need to control synthetic systems. From regulating steam engines to tuning radios to devices capable of autonomous movement, it provided a formal mathematical basis for understanding the role of feedback in the stability (or change) of dynamical systems. It provides a framework for understanding any system with feedback regulation, including biological ones such as regulatory gene networks, cellular metabolic systems, sensorimotor dynamics of moving animals, and even ecological or evolutionary dynamics of organisms and populations. Here we focus on four case studies of the sensorimotor dynamics of animals, each of which involves the application of principles from control theory to probe stability and feedback in an organism's response to perturbations. We use examples from aquatic (electric fish station keeping and jamming avoidance), terrestrial (cockroach wall following) and aerial environments (flight control in moths) to highlight how one can use control theory to understand how feedback mechanisms interact with the physical dynamics of animals to determine their stability and response to sensory inputs and perturbations. Each case study is cast as a control problem with sensory input, neural processing, and motor dynamics, the output of which feeds back to the sensory inputs. Collectively, the interaction of these systems in a closed loop determines the behavior of the entire system.

Keywords: feedback, control theory, neuromechanics, stability, locomotion.

Introduction

The idea that organisms can be understood as a hierarchy of organizational levels—from molecules to behavior—seems intuitive. Indeed, a dominant paradigm in biological science involves a reductionist approach in which a phenomenon at a particular level of organization is described as a consequence of the mechanisms at a lower level. For example, the mechanisms underlying behavior might be described in terms of the activity of a set of neurons, or the behavior of a single neuron might be understood in terms of its membrane properties.

However, each level of organization exhibits emergent properties that are not readily resolved into components (Anderson, 1972), suggesting the need for an integrative approach. The case for taking an integrative view is strengthened by considering a fundamental organizational feature inherent to biological systems: feedback regulation. Living systems ubiquitously exploit regulatory mechanisms for maintaining, controlling, and adjusting parameters across all scales, from single molecules to populations of organisms, from microseconds to years. These regulatory networks can form functional connections within and across multiple levels (Egiazaryan and Sudakov, 2007). This feedback often radically alters the dynamic character of the subsystems that comprise a closed-loop system, rendering unstable systems stable, fragile systems robust, or slow systems fast. Consequently, the properties of individual components (e.g. biomolecules, cells, organs), the communication channels that link them (e.g. chemical, electrical, mechanical), and the signals carried by those channels (e.g. phosphorylation, action potentials, forces), can only be understood in terms of the performance of the complete feedback control system. Understanding the role of individual components in the context of a complete feedback system is the purview of control systems theory (Astrom and Murray, 2008). Control theory provides a suite of tools and language for describing biological feedback control systems (Roth et al., 2014).

An organism comprises a complex patchwork of feedback control systems that cut across traditional levels of biological organization. Thus, understanding biological systems requires an understanding of what feedback can (and cannot) do. Feedback can be used to dramatically enhance robustness and performance of a system. However its benefits are not endless: there are inherent, often inescapable tradeoffs in feedback systems, and control theory provides precise quantitative language to address such tradeoffs (Middleton, 1991; Looze et al., 2010; Freudenberg and Looze, 1985).

Perhaps nowhere are feedback control tradeoffs—such as the intricate balance between stability and change—more immediately relevant than in the controlled movement of animals. While performance measures such as speed and efficiency are essential to some behaviors, some measure of stability almost invariably plays a role, as the fastest animal would have poor locomotor performance if the smallest irregularity in the surrounding environment was sufficient to cause it to crash, fall, break, or fail (Dickinson et al., 2000). Yet animals do not seem to adopt the most stable, conservative designs—

e.g. aggressively regulating convergence rate and bringing the animal to rest as quickly as possible may be inefficient from a different perspective (Ankaralı et al., 2014).

There is no single definition of "stability" or "change" that is ideally suited for all biological systems. In neuromechanical terms, however, "change" can be defined in terms of the responsiveness or maneuverability of the motor system (Sefati et al., 2013) to a sensory stimulus—i.e. the "bandwidth" of the system. "Stability" on the other hand, refers most generally to the ability of a system to reject external perturbations, but can also refer to the "persistance" of a system (Ankaralı et al., 2014; Byl and Tedrake, 2009).

After providing a short historical perspective on feedback control and biology, we review a diversity of sensorimotor feedback control systems. The methods reviewed in this paper are remarkably conserved, despite categorical differences in species (vertebrates and invertebrates) and locomotor modalities (terrestrial, aquatic, and aerial). Indeed, one of the sensorimotor behaviors, the jamming avoidance response in the weakly electric fish, requires no movement, but yet enjoys the same basic modeling tools and approaches as used for the movement based sensorimotor feedback systems.

A Brief History of Feedback Control and Living Systems

In the latter part of the 18th century, James Watt and Matthew Boulton invented an ingenious device that provided feedback control for the steam engine. In this device, modeled after similar mechanisms used in windmills and millstones (Hills, 1996), the passage of steam was controlled by a governor (Figure 1) in which the rotational velocity of the engine's output was "sensed" by two masses suspended from an articulating lever system. As the rotation rate of the engine increased, the centripetal acceleration raised the masses closing the throttle to the engine. Thus the output of the engine was sensed by a physical mechanism that influences the input of steam into the engine. The governor is an elegant physical instantiation of a closed-loop feedback system: the spin-rate of the mass depends on, and controls, the flow of steam through the system.

That concept of a governor—a closed-loop feedback system—has its tendrils in evolution and ecology. Indeed, the idea that feedback plays a central role in evolution owes its origins to Alfred Wallace (1858):

> The action of this principle is exactly like that of the centrifugal governor of the steam engine, which checks and corrects any irregularities almost before they become evident; and in like manner no unbalanced deficiency in the animal kingdom can ever reach any conspicuous magnitude, because it would make itself felt at the very first step, by rendering existence difficult and extinction almost sure soon to follow.

At about the same time that Darwin and Wallace were forging the principles of evolution, the concept of the governor in feedback control extended deeply into dynamical systems theory with James Maxwell's early contribution "On Governors" (1867) in which he lays out the mathematical formulation for the equations of motion of a governor and a speed dependent term throttling down the spin rate. The major contribution of this work was his generalization to a broad class of closed-loop feedback systems, with the novel idea that the input and output of such systems are inextricably linked (as they are in living systems). Maxwell is, in many ways, one of the founding contributors to the theory of control systems.

Following Maxwell, systems and controls theory takes on a rich history with a focus on quantifying the stability and performance of dynamical systems with the seminal contributions of Aleksandr Lyapunov, Harry Nyquist, and Hendrik Bode (among many others) in the late 19th and early 20th centuries. These and subsequent historic contributions of control theory to engineering domains—from tuning radios to aircraft controllers, and much more—are summarized in a short history of the field by Bennett (1996).

The ideas of Wallace and Maxwell can be traced through to living systems in a host of contributions. For example, at the time of Maxwell, the French physiologist Claude Bernard developed the idea (later termed "homeostasis") suggested that physiological systems maintain a constant internal surrounding environment ("la milieu interieur") via physiological feedback. Interestingly, glucose control and its consequences to diabetes was a key example of feedback developed by Bernard. Indeed, this concept of feedback and control is a hallmark of the sort of biological systems we study and teach.

In the latter part of the 19th century and early 20th century, the contributions of dynamical systems theory to living systems began to form with an initial focus on neural systems and cybernetics. The Russian physiologist Pyotr Anokhin for example, developed the idea of "back afferentiation" (feedback) in neural control of sensorimotor systems and reflexes (for a review, see Egiazaryan and Sudakov, 2007). Anokhin's "functional systems theory" has all the hallmarks of dynamical systems and control theory we use in cybernetics and in systems biology today.

While Anokhin was delving into neural systems, reflexes and control, Norbert Wiener, a pioneer in mathematics and engineering, had begun to focus much of his attention on the control laws associated with animals and machines, essentially founding the domain of cybernetics (Wiener, 1948). Along with the contributions of Ludwig von Bertalanffy (1969), whose mathematical models of animal growth are still used today, the fields of cybernetics and von Bertalanffy's "general systems theory" gave rise to the burgeoning fields of systems neuroscience, systems biology, and robotics. Indeed, the sort of integrative biology we outline in this paper is in every sense "systems biology" without the restriction of attention solely to genetic and molecular scales, but with all the requisite mathematical underpinnings that began with Watt and Maxwell.

Four Systems that Walk the Proverbial Tightrope

We turn our focus to the dynamics and control of motor behavior in animal systems (neuromechanics) to illustrate basic



Figure 1: (A) Schematic of the Watt-Boulton centrifugal "flyable" governor (copyright expired; see `http://copyright.cornell.edu/resources/publicdomain.cfm`) and (B) a simplified feedback diagram.

conceptual issues surrounding the application of control theoretic analyses that address the apparent dichotomy between stability and change. Specifically, we provide four examples of control theoretic analyses of neuromechanical systems. For each example, we describe the most important features of the control system at hand, point out any task-relevant tradeoffs, and discuss how the organism walks the proverbial tightrope.

A second but equally important goal is to review the application of control theoretic analyses in interpreting the roles of constituent components of a biological feedback control system. The specific physical details of individual components of the system are most meaningfully described in the context of the intact control circuit. With this goal in mind, we review rapid wall-following in cockroaches, refuge tracking in weakly electric fish, the abdominal reflex response in hawk moths, and the jamming avoidance response, again in weakly electric fish.

Cockroach Wall Following

Central to understanding how animals manage stability and change is evaluating performance in challenging contexts (Dickinson et al., 2000). Crucially, this involves creating significant deviations from steady-state behavior. Moreover, it encourages us to consider organisms and behaviors that have evolved in environments where the pressures, and potential tradeoffs, between stability and change may be finely balanced. Locomotion is particularly challenging when it is difficult to move (surfaces can be irregular and deformable (Daley and Biewener, 2006; Li et al., 2009; Sponberg and Full, 2008; Li et al., 2013), the surrounding fluid can be turbulent and cluttered, or difficult to sense (some highly visual animals navigate in very low light (Warrant and Dacke, 2011)), and others behave in complex auditory (Stamper et al., 2008) or olfactory landscapes. Behaving in such environments can be advantageous to avoid competition, predation, and enable intraspecific interactions.

Despite significant neural resources dedicated to vision, many species of cockroach navigate low-light and cluttered environments (Bell et al., 2007). Rapid locomotion through rough, cluttered or deformable terrain can render vision unreliable, particularly given its longer latencies compared to other senses (Franklin and Wolpert, 2011). Sensor latency can pose fundamental limits on neural control, and delays are an inevitable part of processing information in biological wetware (Cowan et al., 2006; Sponberg and Full, 2008; Elzinga et al., 2012; Ristroph et al., 2013; Fuller et al., 2014). One strategy cockroaches use is to probe their environment using their antennae as tactile sensors, mechanically detecting and following vertical surfaces (or "walls") during rapid running (Figure 2A) (Camhi and Johnson, 1999; Cowan et al., 2006; Lee et al., 2008). Thousands of mechanoreceptive hairs lining the cockroach's antennae are activated on contact and bending (Schafer and Sanchez, 1973; Schaller, 1978; Camhi and Johnson, 1999; Cowan et al., 2006). The mechanics of the antenna itself passively maintains the orientation of the antenna in a "J" shape against the wall (Figure 2A) (Mongeau et al., 2013). The cockroach runs while maintaining close proximity to the wall and tracks turns and irregularities in the surface. This behavior enables remarkably high-bandwidth maneuvers, necessary for maintaining high-speed performance associated with escape and navigation; cockroaches reportedly respond to corrugations in a wall with up to 25 turns per second (Camhi and Johnson, 1999). Blinding the animals does not significantly impair performance and they do not require contact from their body or legs to wall follow (Camhi and Johnson, 1999).

Wall following in cockroaches provides an excellent exam-



Figure 2: (A) Schematic depicting a cockroach following a wall and (B) a simplified block diagram representation of cockroach wall-following behavior. The reference signal is the position of the wall in some global reference, $r(t)$. The difference between the wall and the cockroach's position, $y(t)$, is the error signal, $e(t)$. The error is encoded in antennal mechanoreceptors and transformed by the nervous system, ultimately causing changes in motor commands that act through the animals body dynamics to alter its own position, thereby regulating this feedback error to a desired reference point.

ple of a smooth pursuit or tracking task (Cowan et al., 2006; Lee et al., 2008). An external reference signal, in this case the wall position, is detected by a sensor—the antenna—and the animal's brain and body cooperate to regulate the distance from the wall (Figure 2). A simple ethological description of wall-following is that the animal "tries" to maintain a certain distance from a surface. This qualitative description of the behavior is unfulfilling because it is neither mechanistic—how the animal "tries" is not well understood—nor predictive—we cannot predict how the animal will recover from perturbations or when its performance will degrade. A classic approach in neuroethology would be to identify neurons potentially involved in the behavior and determine what their responses are to a variety of mechanical disturbances. However, the challenge is that many models may explain observed patterns of encoding and the relevant mechanisms may be difficult to identify without rejectable hypotheses derived from quantification of the animal's dynamics. For example, a mechanoreceptive neuron may appear to respond to deflections of the antenna, but its time constants may be too long (or too short) to play a role the wall-following behavior. Control theoretic approaches to understanding the dynamics of both the neural processing and the body's movement enable testable predictions that inform behavioral, neurophysiological, and biomechanics experiments.

The control theoretic approach characterizes wall following as a feedback-regulated behavior (Figure 2B). Using biomechanics models for the body dynamics—either stride-averaged (Cowan et al., 2006) or continuous (Lee et al., 2008))—we can implement different hypothesized models for how the nervous system processes the error signal (i.e. the mechanical bending of the antenna) and compare the resulting dynamics of the whole feedback system to kinematics of cockroaches following turns in walls. This has led to the discovery that the animal must encode more than just the position of the wall. The simplest control model that matches behavior requires the position and rate of approach of the wall (i.e. the lateral velocity of the wall relative to the cockroach) (Cowan et al., 2006). Such a PD (position and derivative) controller is ubiquitous in controlled engineering systems. However, while the latency of the initial turn is low, the turning response persists for much longer than the stimulus (Lee et al., 2008). The system dynamics filter the sensory input in time. Electrophysiology of the antennal nerve revealed that the population of mechanoreceptors in the antennal nerve could encode both position and velocity and that the neural response even at this first stage of processing is already temporally matched to the turning response rather than to the stimulus (Lee et al., 2008). In other words, the mechanosensory processing seems to be tuned to the demands imposed on the control system by the mechanics and the neural processing delay. A hallmark of a high-performance control system is the ability to achieve large responses over a wide frequency range in response to stimuli (change) without skirting too close to the instabilities that can result from high-gain, large-latency feedback (stability).

Refuge tracking in weakly electric fish

The glass knifefish, *Eigenmannia virescens*, is like an "aquatic hummingbird": it hovers in place with extraordinary precision, making rapid and nuanced adjustments to its position in response to moving stimuli (Roth et al., 2011; Cowan and Fortune, 2007; Rose and Canfield, 1993a,b); see Figure 3A. Here, we investigate the integration of locomotor biomechanics (Sefati et al., 2013) (Figure 3B), multisensory intgration (Stamper et al., 2012b), and adaptive control (Roth et al., 2011) that enable this animal to balance stability and change. See Figure 3.

As a model organism, weakly electric knifefish are most widely studied for their namesake, an active electrosensory system. An electric organ (EO) in the tail emits a weak electric field. Electroreceptors distributed over the surface of the body (most densely about the head) detect objects in the near field as small disturbances in transdermal potential. Using this electrosense in conjunction with vision, fish perform a wide variety of localization and tracking behaviors. As in the analyses presented for the wall-following behavior in cockroaches, we again demonstrate how a control theoretic approach can be used to quantify and model behavior. For the fish, we further extend the modeling tool to quantitatively probe the categorical shifts in behavior and the interplay between visual and electrosensory modalities (Figure 3C).

Weakly electric knifefish hunt nocturnally, their specialized electrosensory system allowing them to navigate their environment and localize small prey items in low light (MacIver et al., 2001). During the day, they hide from predators, find-



Figure 3: (A) The knifefish in a moving shuttle. Positions are measured from a fixed reference frame to tracking points on the refuge and the animal body. (B) A schematic depicting the counter-propagating wave kinematics of the knifefish ribbon fin. As ambient flow velocity, u, increases, fish recruit a larger portion of the fin for L_{head}, the wave component responsible for forward thrust. (C) A block diagram depicting the knifefish reference tracking behavior. The moving shuttle provides the reference signal, $r(t)$, with the output, $y(t)$, being the position of the swimming fish. Parallel visual and electrosensory modalities measure the relative position of the shuttle (the sensory slip, $e(t)$). The neural controller (CNS) weights and filters signals from the sensory blocks and outputs commensurate motor commands. Subsequently, these motor commands generate movement as dictated the biomechanics of the fish body and the body-fluid interaction.

ing refuge in tree root systems or other natural shelter. In the laboratory, these fish exhibit a similar refuge-seeking behavior, hiding in short lengths of pipe, filter fixtures or any other refuge provided for them. More impressively, fish smoothly and robustly track their refuge as it is moved (Figure 3A). How do sensorimotor control strategies differ across this repertoire, in response to different categories of exogenous motion? And how do visual and electrosensory cues contribute to these behaviors?

Knifefish are agile. An undulating ribbon-fin runs along the underside of the body, enabling knifefish to rapidly alternate between forward and backward swimming without changing body orientation. In experiments, this remarkable ability is often leveraged to constrain the behavior to a line of motion, reducing the spatial dimension of the task to single degree of freedom. Fish were first observed swimming side-to-side in response to laterally moving plates and rods, termed the "following" response (Heiligenberg, 1973b; Matsubara and Heiligenberg, 1978; Bastian, 1987b) and later experiments explored similar behavior in response to longitudinally (fore–aft) moving refuges (Rose and Canfield, 1993a, 1991; Cowan and Fortune, 2007; Roth et al., 2011; Stamper et al., 2012b). In the case of longitudinal refuge tracking, the fish maintains a "goal" position within the refuge, perceiving the error between its position and that of the refuge and swimming forward or backward to minimize this positional error (Figure 3A and C).

Linear control theoretic tools were used to characterize the (frequency-dependent) relationship between sensing and swimming in this smooth-pursuit behavior (Cowan and Fortune, 2007). Interestingly, however, subsequent investigations revealed important deviations from the linear model proposed in (Cowan and Fortune, 2007). When comparing the responses to complex motion (in this case, sums of sinusoidal trajectories across a broad band of frequencies) and pure sinusoidal stimuli, fish exhibited performance suggesting qualitatively different underlying models (Roth et al., 2011). This failure of the so-called superposition property of linear systems, revealed an interesting nonlinearity: fish tune their control policy to improve behavior with respect to the spectral content of the stimulus. While this adaptive tuning improves the response to a limited class of stimuli (and these may well be of greater ethological relevance), performance suffers in response to signals that contain broader frequency content.

Another tradeoff manifests as a nonlinearity in the context of active sensing. When visual cues are limited in the environment (e.g. in low illumination or murky waters) fish rely predominantly on electrosensation. Under such circumstances, a conflict arises between the goal of the tracking task—to remain stationary within the tube—and the needs of the electrosensory system—which requires active movements to prevent adapting to a constant sensory stimulus. In low light conditions, the animal performs a rapid back and forth shimmy superimposed on the tracking response. These volitional motions are uncorrelated to the refuge motion and can be discriminated from the tracking response by their frequency content. While these active oscillations are significant, tracking performance with respect to the stimulus motion remains nearly constant (Stamper et al., 2012b). It is posited that these oscillations serve to enhance electrosensory acuity and permit a high level of performance in the absence of salient visual stimuli. Fish employ a strategy in which tracking error and expended energy are compromised for improved sensing and tracking of the stimulus.

While we present the tracking behavior as a model system for the study of smooth pursuit and sensory integration, the locomotor strategy also illustrates a behavioral tradeoff. Knifefish routinely partition the ribbon-fin into two counter-propagating waves (Sefati et al., 2010; Ruiz-Torres et al., 2013), recruiting the frontal portion of the fin to generate forward thrust (a head-to-tail traveling wave) with the rear section (tail-to-head wave) generating an opposing force. In

stationary hovering, these opposing forces cancel each other. These waves meet at the "nodal point" (Figure 3B); simulations on a biomimetic robot reveal that the net fore-aft thrust is linearly dependent on this kinematic parameter(Sefati et al., 2013; Curet et al., 2011). Moreover, when compared to simpler strategies (e.g. recruiting the whole fin and modulating the speed of a single traveling wave), the use of counter-propagating waves significantly improves the fore–aft maneuverability (by decreasing the control effort) and concurrently enhances the passive stability (stabilization without active feedback control) by providing a damping-like force to reject perturbations, thus simplifying the neural control.

Moth Abdominal Reflex For Pitch Control

Here, we examine the problem of active pitch control in the hawk moth, *Manduca sexta*, as a platform to explore the relationship between open-loop experiments and closed-loop stability and maneuverability. The inherent instability of flapping flight requires active, feedback-based strategies for control (Wu and Sun, 2012; Liang and Sun, 2013). This has led to the evolution of numerous sensory specializations, most notably in the form of visual and mechanical senses (O'Carroll et al., 1996; Pringle, 1948; Sane et al., 2007; Taylor and Krapp, 2007), that collectively inform the animal about its state in the environment. This information, in turn, is used to coordinate motor systems to direct movement (Figure 4).

The bulk of research on animal flight control has focused on how the wings are used to generate and modulate aerodynamic forces. Much less attention has been paid to the role of body—or "airframe"—deformations for flight control. The hawk moth displays strong abdominal movements in response to open-loop visual rotations during tethered flight. Control theory can provide key insights about the importance, and possible advantages, of such movements for controlling flight.

Numerous experimental preparations have been developed for investigating sensory and motor responses that involve restraining or confining animals to access physiological signals and to allow for better experimental control of the sensory inputs available to the animal. For hawk moths, these include a tethered virtual flight arena for performing behavioral experiments (Figure 4A) to immobilized and dissected preparations for electrophysiological recordings (Hinterwirth and Daniel, 2010; Theobald et al., 2010). However, these types of manipulations dramatically change the dynamic context of the animal. The difficulty then is linking physiological signals and behavioral responses from restrained preparations to free flight movements, such that causal links can be made between sensing, movement and flight path changes.

It is here that the analytic techniques of control theory provide unique affordances for understanding how animals control flight. Control theory provides a framework for interpreting data from restrained experimental preparations in the context of the free flight dynamics via mathematically explicit dynamical models derived from the basic physics and mechanics of flight. In turn, these studies provide movement predictions that can be compared to natural flight behaviors.

Hawk moths are accomplished fliers and spend much of their time during flight hovering in front of flowers while feeding. Hovering flight is an equilibrium flight mode that makes the modeling particularly tractable for control theoretic analyses. Dyhr et al. (2013) took advantage of these simplified dynamics to test the utility of abdominal responses for pitch stabilization. Previous studies had suggested that abdominal movements were of minor importance for flight control (Hedrick and Daniel, 2006; Cheng et al., 2011). But, absent a control theoretic analysis of free flight, it is difficult to exclude a crucial role for the deformation of the airframe (redistribution of mass).

Free flight is a closed-loop behavior such that movements the animals make influence subsequent stimuli. Restraining an animal so that open-loop responses—in which animal movements no longer influence the sensory input—can be measured and often generate stronger responses and more data, simplifying the quantification of the sensorimotor transform. However, these data can only be interpreted in their closed-loop context (Roth et al., 2014). Dyhr et al. (2013) combined behavioral experiments, in the form of open-loop tethered flight responses, with a hovering flight dynamics model to generate a control theoretic model of closed-loop visual-abdominal control. See Figure 4. Using this model they were able to show that visually evoked abdominal movements were sufficient for pitch stabilization during hovering flight independent of any modulation or redirection of the wing forces.

While this work has demonstrated the importance of airframe deformation for flight control, other actuator systems are clearly involved. The wings are certainly the most important structures for flight control, but head movements have an established role in modulating both visual and mechanosensory information (Dyhr et al., 2013; Hinterwirth and Daniel, 2010). Understanding how wing, body, and head movements are coordinated is a promising future research direction. Future work in this area will require integrating realistic aerodynamic models with rigid body dynamics to understand how multi-input control is achieved. This problem is also exciting from a multisensory integration standpoint, as hawk moths use multiple sensory modalities (e.g. visual and antennal mechanosensory, Sane et al., 2007; Hinterwirth and Daniel, 2010) for flight control. Furthermore, investigations into the coordinating multiple motor pathways may highlight the importance of proprioceptive feedback mechanisms for flight control, an area that has been relatively unexplored. The tractability of *Manduca sexta* as an experimental organism for both behavioral and physiological studies, coupled with the relatively simple dynamics of hovering flight, make it a promising model organism for understanding the sensorimotor processes underlying locomotor control.

Jamming Avoidance

As the previous examples show, control theory is a useful tool for understanding sensorimotor systems, especially during behaviors which are robust and repeatable. However, escape responses represent a behavioral category where the animal actively tries to avoid a particular sensory condition. From



Figure 4: (A) Experimental setup for measuring responses to visual pitch perturbations in *M. sexta*. The moth is attached to a rigid tether and placed in a cylindrical LED arena. During bouts of flight the moth is presented with either an isolated visual stimulus, $r_1(t)$, by rotating a green and black striped pattern on the visual display, an isolated mechanical stimulus, $r_2(t)$ by physically rotating the moth and the arena, or a coupled visual and mechanical rotation. The moth responds to the rotations by moving its head (blue), wings (red) and abdomen (yellow, $y(t)$). (B) Block diagram of the different sensory and motor system known to be engaged during open-loop tethered flight. Error signals, $e_i(t)$, represent perceived visual (eyes) and mechanical (antennae) sensory information relative to environmental reference signals, $r_i(t)$. Sensory systems independently encode the modality specific signals and are then fused and processed by the nervous system. Neural commands are relayed to different motor systems to achieve new kinematic states. (C) Block diagram combining open-loop experimental data (visual-abdominal transfer function) and dynamics models (mechanical model) to estimate the behavior of the closed-loop system (Dyhr et al., 2013).

a control systems perspective, the behavior is transient: it "escapes" to the nearest stable equilibrium. The response is not amenable to perturbation analyses that we have used so far, and modeling such a response requires a different approach, such as experimentally "closing" the loop. Here, we re-examine weakly electric fish in the context of just such an unstable sensorimotor escape behavior.

In addition to sensing the environment for behaviors such as tracking, described above, the electric organ discharge (EOD) is used for social communication. In wave-type fish, each individual produces a continuous, pseudo-sinusoidal EOD whose frequency and amplitude can remain remarkably constant for many hours and even days (Bullock et al., 1972). However, if two nearby fish have frequencies F_1 and F_2, their electric fields interact to produce a "beat" at the difference frequency $|\mathrm{dF}| = |F_1 - F_2|$. When the two frequencies are within a few Hz of each other, the emergent low-frequency beat detrimentally interferes with electroreception, thereby "jamming" the ability of the fish to detect obstacles and prey (Heiligenberg, 1973a; Bastian, 1987a). Some species of these fish, particularly those that form social groups (Stamper et al., 2010), can rapidly change their frequencies to avoid such interference. This behavior is termed the Jamming Avoidance Response (JAR).

The neural computation that underlies the JAR in the glass knifefish, *Eigenmannia virescens*, has been elucidated via a half century of research (Heiligenberg, 1991; Fortune, 2006). The JAR can be predicted on the basis of a single parameter, the dF which can be used to predict the structure of electrosensory beats. Using parallel receptor systems, the fish encodes amplitude modulations and relative phase modulations of zero crossings in the beats to drive a motor response that serves to increase the magnitude of the dF.

But this is not the whole story. Each individual tends to return to its own internal EOD frequency set point in the absence of a low |dF|. The set point can drift over long periods of time, but over the timescales of the JAR (seconds to minutes) the EOD set point remains constant. This return to baseline likely serves to maintain the EOD within the range of frequencies that match the tuning properties of each individual's own electrosensory receptors. For example, if a fish were producing a 400 Hz EOD, its receptors would be most responsive to EOD frequencies within a range of approximately 300 to 500 Hz (Scheich et al., 1973).

How do individuals balance the need for maintaining electrosensory stability while still being able to rapidly change the EOD frequency during the JAR? To address this question, Madhav et al. (2013) modeled the JAR in terms of a low-order feedback control system that includes both the return to baseline (stability) and the stimulus-driven escape (change). Parsing this feedback control diagram (Figure 5) requires understanding that the JAR operates on frequencies of signals. Specifically, the system computes the instantaneous difference between an exogenous input (conspecific frequency) and the autogenous output (self frequency) as shown in Figure 5. This difference is dynamically processed by the CNS which in turn



Figure 5: (A) Experimental setup to identify the JAR in *Eigenmannia*. The fish is placed in a tube in the experimental tank, and recording electrodes (red) are used to measure its EOD. The EOD is amplified and its frequency is extracted. This frequency is fed to the controller which generates the appropriate input frequency based on a control law. A signal generator outputs a sinusoid at the input frequency, which is then played into the tank through the stimulus electrodes (black), through a stimulus isolation unit (SIU). (B) Block diagram representation of the same experimental paradigm. The reference, $r(t)$, output, $y(t)$, input, $u(t)$, and dF, $d(t)$, are all frequency signals relative to the baseline frequency of the fish. We seek to identify the unstable open loop (green dashed box) using the stabilized closed loop (orange dashed box). The dF computation is modeled to have a lumped delay. The delayed difference initiates the sensory escape, which competes with the motor return to produce the output EOD frequency.

modulates the output, creating a closed feedback loop.

The challenge in identifying the dynamics of the JAR was twofold. First, the tendency of the output to diverge from the input renders the system locally unstable. Effective system identification techniques rely on analyzing persistent responses to perturbations; however, in this case, these perturbations destabilize the system, making it impossible to apply such techniques. However, this challenge was overcome by stabilizing the behavior using an experimentally closed loop (Figure 5). This stable closed-loop system was systematically perturbed and the responses were used to identify a linear model, which describes the unstable open-loop behavior in the local neighborhood of the baseline frequency of the fish. Second, predictions of responses in real-world scenarios requires understanding the global nonlinear nature of the behavior. A different category of closed-loop experiments were used to identify a characteristic "escape curve", which serves as the nonlinear signature for the JAR for each individual. Identifying this curve for each individual allows us to populate all the parameters of the global model.

In the global model, the computational algorithm of the JAR was expressed as a competition between the stable motor dynamics (return to baseline) and the need to adapt to changing social settings (sensory escape). Comparatively simple behavioral experiments can now be used to fit parameters of this model, which can in turn predict responses to naturalistic or novel artificial stimuli. For example, this model captures the asymmetry between rises and falls in EOD frequency, for which a neural correlate was described previously (Metzner, 1993). This model could also be used to predict social interactions between two individuals without considering the details of each individual's behavioral characteristics. This model can also form a basis for future work investigating complex social interactions of three or more individuals, where higher-order electrosensory envelopes can also drive behavior (Stamper et al., 2012a).

Discussion

We have seen from the above examples that control theory and system identification tools give us a quantitative framework in which to interpret comparative organismal studies of locomotion. In cockroach wall following, we tested hypotheses about neural encoding derived from the sufficiency of simple control laws and mechanical models. In the fish swimming and moth flying examples, we understood the contributions of multiple sensory signals and multiple actuators (a.k.a. multiple-input, multiple-output or MIMO) to the production of movement. Along the way we discovered new principles about how animals are dynamically tuned to their environment. This perspective allows us to relate the results of closed and open-loop experiments and test hypothesis about stability and maneuverability. With the jamming avoidance response, we saw how control theory applies to behaviors that move not in physical space, but in the frequency space animals use for infraspecific signaling and communication. These questions of stability, change, sensing and movement are fundamental to the emerging field of systems neuromechanics. We can use the language of control theory to translate between open- and closed-loop experimental preparations (Roth et al., 2014), allowing us to relate the functional mechanisms of individual components to the integrated performance of the intact, behaving animal, just as the early pioneers, such as Bernard, Anokhin, and Wiener, envisioned.

Closing the Loop from Biology to Control Theory

Just as control theory affords rich insight into the role of stability and change in living systems, there is a feedback loop that couples research on living systems back to the tools we need from control theory. It is crucial to realize that while control theoretic tools can enable biologists to tackle challenging problems of great significance (biomedical, evolutionary, and environmental) the same can be said of the impact of biology on development of tools in engineering and control

theory. We recall that in fields such as physics and fluid dynamics, the need for models and mathematical formalisms continues to spur the development of powerful computational and analytic methods.

Because biological phenomena are much more complex—chemically, physically, and organizationally—than inorganic phenomena, a cause-and-effect understanding of such complex systems will inevitably foster innovative analytic, computational and technological advances. Some key examples emerging today include the need for new analytic methods estimating the dynamics of freely behaving animals (Revzen and Guckenheimer, 2011) and reverse engineering biological networks (Kang et al., 2013).

Integrating Empirical and Physics-Based Models

The nervous system processes the sensory information for closed-loop control of task-level locomotion, such as tracking behavior (Cowan and Fortune, 2007; Rose and Canfield, 1993b). In control systems terminology, the mechanical *plant* defines the way motor signals are transformed into forces and movements, and so discovering the neural controller (Ekeberg, 1993; Frye and Dickinson, 2001; Nishikawa et al., 2007; Roth et al., 2011; Miller et al., 2012) of a biological system greatly benefits from a task-specific mechanical model of the underlying locomotor dynamics (Sefati et al., 2013; Cowan and Fortune, 2007; Cowan et al., 2006). Low-dimensional, task-specific models for the locomotor mechanics enable the application of control systems analysis to understand the neural mechanisms for sensorimotor processing (Blickhan and Full, 1993; Holmes et al., 2006; Cowan et al., 2006; Hedrick and Robinson, 2010; Tytell et al., 2011). These simple descriptive mechanical models, sometimes termed "templates" (Holmes et al., 2006; Full and Koditschek, 1999), are essential for understanding stability and control in biological systems (Blickhan and Full, 1993; Sefati et al., 2013; Schmitt and Holmes, 2000).

More elaborate models, sometimes termed "anchors" (Holmes et al., 2006; Full and Koditschek, 1999) can facilitate the exploration of more detailed questions about closed-loop control. Multidisciplinary approaches integrate computational models and experiments with biomimetic robots to study the locomotor mechanics in more details and with higher accuracy Miller et al. (2012). With advances in computing, high-fidelity simulations have categorically improved our understanding of various locomotor strategies in different species (Mittal, 2004; Wang et al., 2004; Luo et al., 2008; Shirgaonkar et al., 2008; Tytell et al., 2010). On the other hand, biomimetic robots enable us to experimentally validate the mechanical models (Wang et al., 2004; Lauder et al., 2007; Sefati et al., 2012, 2013), and to explore the effect of parameters beyond their biological ranges, providing insight as to where the biological performance lies within the range of the wider range of possible mechanical solutions (Curet et al., 2011; Sefati et al., 2013).

Neurophysiology

How can the insights concerning the role of feedback in the maintenance of stability and the control of change at the organismal level be used to decode neurophysiological mechanisms used in the brains of animals? First we need to determine what we want to learn from and about the nervous system. In terms of whole animal control, the nervous system is simply one part of the closed-loop system. In this context, understanding the inputs and outputs of the nervous system under behaviorally relevant conditions might be sufficient. The nervous system can remain a black box that is used to better understand the behavior at the organismal level in its native closed-loop state. Alternatively, we may use insights from behavior as a tool for understanding the properties of the nervous system as a functional unit. Indeed, studying the nervous system within a closed-loop behavioral task is perhaps the best route for understanding the functional structure and organization of the nervous system. In this case, behavior is used to understand the sets of computations within the black box.

The central challenge in decoding neurophysiological mechanisms is that brains are typically composed of millions of independent neurons each of which may have unique structure and function. Organism-relevant computations for both maintaining stability and controlling change are often distributed over thousands to millions of neurons that act in parallel. Presently we do not understand the nature of the coding systems that are used in single neurons, and it is unclear what sorts of dimensional reduction are possible across populations and networks of neurons. In other words, there appears to be no simple or obvious set of *a priori* constraints that control theory can contribute to decoding the neurophysiological activity of neurons in the brain. This problem is familiar to neuroscientists, as one of the long standing challenges in the study of neural mechanisms is discovering strategies that effectively translate behavioral observations into feasible neurophysiological experiments.

This challenge stems in part from the fact that neurons are on the order of microns to tens of microns in diameter and use tiny electrical signals. As a result, the vast majority of neurophysiological experiments have relied on the placement of microelectrodes into anesthetized and/or immobilized animals or into neural tissues that have been removed from the animal. Obviously, the critical organism-level feedback systems that are essential for the control of behavior are disrupted in these sorts of experimental preparations. In other words, studies have been conducted in neural tissues in which the closed-loop control system has been opened by the elimination of feedback. This is important because it is almost certainly not possible to extrapolate the neural signals from data obtained in immobilized animals to make control predictions in the intact behaving animal (a notable exception is electric fish, in which certain electrosensory behaviors remain intact in immobilized individuals (Fortune, 2006)). These studies in open-loop preparations cannot capture the dynamics of the closed-loop system, and further, are likely to be misleading (Szwed et al., 2003; Cowan and Fortune, 2007; Roth et al.,

2014).

Thankfully, improvements in neurophysiological techniques are now permitting the widespread recording of neurophysiological activity in the central nervous systems of awake, behaving animals (Nicolelis, 2008) and with stunningly compact wireless and battery less technologies coming to play a greater (Thomas et al., 2012). Similarly, recently developed genetic and optogenetic manipulations (Boyden et al., 2005; Zhang et al., 2007) can also be used in animals in which the behavioral control loop remains intact.

Characterizing and manipulating internal signals during movement

One of the strengths of control theoretic approaches is that we can characterize the relationship between any two signals (neural, muscular, mechanical, etc.) with the organism as a function of the underlying neuromechanical system. When we do gain physiological access we can use the same techniques to relates neural spiking to movement and sensory feedback to muscle activation. While the examples in this paper emphasized monitoring motor output while manipulating a sensory reference single, these signals do not need to be an external input leading to a kinematic output. Direct alteration of feedback, either through dynamic manipulation of sensory feedback or by applying perturbations directly to the constituent neural and mechanical systems during closed- or open-loop behavior, is the hallmark of the control systems approach, but is among the least explored experimental approaches at present (Roth et al., 2014). The ability to inject both noise and alter neural processing during behavior affords separation of the contributions of sensors, controllers, and body dynamics to behavior.

One place where this approach has been used is to identify the role of individual muscles in the control of movement during posture control, running, and flight (Sponberg et al., 2011c; Sponberg and Daniel, 2012). During restrained or free behaviors these experiments precise altered or "rewrote" the activation patterns of individual muscles to identify their role in shaping motor output. In systems where the time constant of the dynamic response to perturbed motor commands is faster than the inherent delay transmitting sensory feedback, this also enables the characterization of open-loop plant dynamics during free behavior (Sponberg et al., 2011b). Alternatively, one can use an open-loop characterization of the mechanics portion of the systems dynamics. By replicating the same patterns of input to the muscle but in a isolated open-loop preparation we can measure the muscles work output (a "workloop") (Sponberg et al., 2011a). Other approaches are beginning to couple environment forces to *in vitro* muscle function via artificially closing a feedback loop between a robotic model and a physiological preparation (Richards, 2011; Richards and Clemente, 2012). From a control theory perspective the classic *in vitro* experimental approaches of neuroscience, muscle physiology, and biomechanics are simply ways to characterize subsystems of the animal (its neurons, muscles, and body–environment respectively) and each result can be synthesized, in explicitly quantitative and mechanistic way, back into an understanding of the dynamics of behavior.

Feedback Control in Biological Systems

Several of the articles in this Special Issue highlight the role of feedback regulation in the apparent dichotomy between stability and change across levels of biological organization, from molecular to ecological. For example, Grünbaum and Padilla (2014) show how ecological demands can trigger phenotypic changes with complex temporal dynamics. Variations in the trophic environment triggers a switch between two phenotypes of radula ("teeth"), ultimately creating a history dependent pipeline of radulae. From the perspective of control theory, these temporal dynamics may create a finite-impulse response filter, allowing the animal to be sensitive to newly available resources (i.e. facilitating change) while maintaining a memory of recently available resources (stability).

At the cellular level, feedback regulation of ATP/ADP is thought to maintain energy homeostasis, but these homeostatic metabolic systems may also regulate the development of the respiratory structures and metabolic pathways that supply oxygen and carbon substrates for energy metabolism. Greenlee et al. (2014) discuss how the development of insect larval tracheal and metabolic systems appear to both sustain metabolic performance and plasticity in the dynamic developmental environment. Such regulation must not, of course, imperil the longer-term developmental outcomes of organisms (Hale, 2014): if developmental processes are too responsive to the environment, they could potentially have deleterious effects on adult structure and performance.

One way to resolve this compromise between stable outcomes and responsiveness to changing resources may be to incorporate a combination of feedback and feedforward control. Indeed, this may help mitigate the tradeoff between preprogrammed developmental cascades—which may be able to produce consistent outcomes, but are unresponsive to environmental demands—and tight feedback regulation—which, while responsive to the environment, may introduce long-term inefficiencies. While fault tolerance is a hallmark of feedback control, even complex feedback control systems are sensitive to certain categories of failure (Csete and Doyle, 2002); such failures in regulatory networks manifest themselves as disease (Nijhout and Reed, 2014). In this way, understanding the mechanisms for feedback regulation in the context of control theory may be a critical step in the treatment of certain diseases. This approach will require the development of new quantitative tools, such as network inference of gene regulatory processes (Ciaccio et al., 2014). When such feedback systems are analyzed using control theory it may enable us to formalize our understanding of the processes that allow biological systems to walk the tightrope between stability and change.

Acknowledgements

This material is based upon work supported by the following grants: The National Science Foundation under grant nos. CISE-0845749 (N. Cowan), IOS-0817918 (E. Fortune and N. Cowan), IOS-0543985 (N. Cowan and E. For-

tune), BCS-1230493 (N. Cowan), IOS-1243801 (D. Padilla), and Postdoctoral Fellowship nos. 1103768 (J. Dyhr); The James S. McDonnell Foundation Complex Systems Scholar Award (N. Cowan); The Office of Naval Research under grants N000140910531 (N. Cowan and E. Fortune) and N000141110525 (N. Cowan); The Komen Endowed Chair (T. Daniel); The Office of Naval Research grant nos. N0014-10-0952 (T. Daniel); The Air Force Research Lab grant nos. FA8651-13-1-004 (T. Daniel); The Air Force Office of Scientific Research grant nos. FA9550-11-1-0155-MODPO4 (T. Daniel).

References

P. W. Anderson. More is different. *Science*, pages 393–396, 1972.

M. M. Ankaralı, H. T. Şen, A. De, A. M. Okamura, and N. J. Cowan. Haptic feedback enhances rhythmic motor control by reducing variability, not improving convergence rate. *J Neurophysiol*, 2014. In press.

K. Astrom and R. Murray. *Feedback systems: an introduction for scientists and engineers*. Princeton university press, 2008.

J. Bastian. Electrolocation in the presence of jamming signals: behavior. *J Comp Physiol A*, 161(6):811–824, 1987a.

J. Bastian. Electrolocation in the presence of jamming signals: Behavior. *J Comp Physiol A*, 161:811–824, 1987b.

W. J. Bell, L. M. Roth, and C. A. Nalepa. *Cockroaches: ecology, behavior, and natural history*. JHU Press, 2007.

S. Bennett. A brief history of automatic control. *IEEE Contr Syst Mag*, 16(3):17–25, 1996.

R. Blickhan and R. J. Full. Similarity in multilegged locomotion: Bouncing like a monopode. *J Comp Physiol A*, 173(5):509–517, Nov. 1993.

E. S. Boyden, F. Zhang, E. Bamberg, G. Nagel, and K. Deisseroth. Millisecond-timescale, genetically targeted optical control of neural activity. *Nature Neurosci*, 8(9):1263–1268, Sep 2005.

T. H. Bullock, R. H. Hamstra, and H. Scheich. The jamming avoidance response of high frequency electric fish. *J Comp Physiol A*, 77(1):1–22, 03 1972.

K. Byl and R. Tedrake. Metastable walking machines. *Int J Robot Res*, 28(8):1040–1064, 2009.

J. M. Camhi and E. N. Johnson. High-frequency steering maneuvers mediated by tactile cues: antenna wall-following in the cockroach. *J Exp Biol*, 202:631–643, 1999.

B. Cheng, X. Deng, and T. L. Hedrick. The mechanics and control of pitching manoeuvres in a freely flying hawkmoth (Manduca sexta). *J Exp Biol*, 214(24):4092–4106, Dec. 2011.

M. F. Ciaccio, J. D. Finkle, A. Y. Xue, and B. N. A systems approach to integrative biology: elucidating association, architecture, and regulation. *Integr Comp Biol*, 2014. In review.

N. J. Cowan and E. S. Fortune. The critical role of locomotion mechanics in decoding sensory systems. *J Neurosci*, 27(5):1123–1128, 2007.

N. J. Cowan, J. Lee, and R. J. Full. Task-level control of rapid wall following in the American cockroach. *J Exp Biol*, 209(9):1617–1629, 2006.

M. E. Csete and J. C. Doyle. Reverse engineering of biological complexity. *Science*, 295(5560):1664–1669, Mar. 2002.

O. M. Curet, N. A. Patankar, G. V. Lauder, and M. A. MacIver. Mechanical properties of a bio-inspired robotic knifefish with an undulatory propulsor. *Bioinspiration & Biomimetics*, 6(2):026004, 2011.

M. A. Daley and A. A. Biewener. Running over rough terrain reveals limb control for intrinsic stability. *Proc Nat Acad Sci*, 103(42):15681–15686, Oct. 2006.

M. H. Dickinson, C. T. Farley, R. J. Full, M. A. R. Koehl, R. Kram, and S. Lehman. How animals move: An integrative view. *Science*, 288(5463):100–106, 2000.

J. P. Dyhr, T. L. Daniel, K. A. Morgansen, and N. J. Cowan. Flexible strategies for flight control: an active role for the abdomen. *J Exp Biol*, 216(9):1523–1536, 2013.

G. G. Egiazaryan and K. V. Sudakov. Theory of functional systems in the scientific school of P.K. Anokhin. *J Hist Neurosci*, 16(1-2):194–205, 2007.

Ö. Ekeberg. A combined neuronal and mechanical model of fish swimming. *Biol Cybern*, 69(5-6):363–374, 1993.

M. J. Elzinga, W. B. Dickson, and M. H. Dickinson. The influence of sensory delay on the yaw dynamics of a flapping insect. *J R Soc Interface*, 9(72):1685–1696, 2012.

E. S. Fortune. The decoding of electrosensory systems. *Curr Opin Neurobiol*, 16(4):474–80, Aug. 2006.

D. W. Franklin and D. M. Wolpert. Computational mechanisms of sensorimotor control. *Neuron*, 72(3):425–442, 2011.

J. S. Freudenberg and D. P. Looze. Right half plane poles and zeros and design tradeoffs in feedback systems. *IEEE Trans Autom Control*, 30(6):555–565, 1985.

M. Frye and M. Dickinson. Fly flight, a model for the neural control of complex behavior. *Neuron*, 32(3):385–388, Nov. 2001.

R. J. Full and D. E. Koditschek. Templates and anchors: Neuromechanical hypotheses of legged locomotion. *J Exp Biol*, 202:3325–3332, 1999.

S. B. Fuller, A. D. Straw, M. Y. Peek, R. M. Murray, and M. H. Dickinson. Flying drosophila stabilize their vision-based velocity controller by sensing wind with their antennae. *Proc Nat Acad Sci*, 2014. In press.

Greenlee, Montooth, and Helm. Predicting performance and plasticity in the development of respiratory structures and metabolic systems. integrative and comparative biology. *Integr Comp Biol*, 2014. In review.

D. Grünbaum and D. Padilla. An integrated modeling approach to assessing linkages between environment, organism and phenotypic plasticity. *Integr Comp Biol*, 2014. In review.

M. E. Hale. Stability and change in the development of movement systems: A case study of pectoral fin development. *Integr Comp Biol*, 2014. In review.

T. L. Hedrick and T. L. Daniel. Flight control in the hawkmoth Manduca sexta: the inverse problem of hovering. *J Exp Biol*, 209(16):3114–3130, Aug. 2006.

T. L. Hedrick and A. K. Robinson. Within-wingbeat damping: dynamics of continuous free-flight yaw turns in Manduca sexta. *Biol Letters*, 2010.

W. Heiligenberg. Electrolocation of objects in the electric fish *Eigenmannia* (rhamphichthyidae, gymnotoidei). *J Comp Physiol*, 87(2):137–164, 1973a.

W. Heiligenberg. "electromotor" response in the electric fish *Eigenmannia* (Rhamphichthyidae, Gymnotoidei). *Nature*, 243:301–302, 1973b.

W. Heiligenberg. *Neural nets in electric fish*. The MIT Press, 1991.

R. L. Hills. *Power from wind: a history of windmill technology*. Cambridge University Press, 1996.

A. J. Hinterwirth and T. L. Daniel. Antennae in the hawkmoth Manduca sexta (Lepidoptera, Sphingidae) mediate abdominal flexion in response to mechanical stimuli. *J Comp Physiol A*, 196(12):947–956, Sept. 2010.

P. Holmes, R. J. Full, D. Koditschek, and J. Guckenheimer. The dynamics of legged locomotion: Models, analyses, and challenges. *SIAM Review*, 48(2):207–304, 2006.

T. Kang, J. T. White, Z. Xie, Y. Benenson, E. Sontag, and L. Bleris. Reverse engineering validation using a benchmark synthetic gene circuit in human cells. *ACS synthetic biology*, 2 (5):255–262, 2013.

G. V. Lauder, E. J. Anderson, J. Tangorra, and P. G. Madden. Fish biorobotics: kinematics and hydrodynamics of self-propulsion. *J Exp Biol*, 210(16):2767–2780, 2007.

J. Lee, S. N. Sponberg, O. Y. Loh, A. G. Lamperski, R. J. Full, and N. J. Cowan. Templates and anchors for antenna-based wall following in cockroaches and robots. *IEEE Trans Robot*, 24(1): 130–143, Feb. 2008.

C. Li, P. B. Umbanhowar, H. Komsuoglu, D. E. Koditschek, and D. I. Goldman. Sensitive dependence of the motion of a legged robot on granular media. *Proc Nat Acad Sci*, 106(9):3029–3034, 2009.

C. Li, T. Zhang, and D. I. Goldman. A terradynamics of legged locomotion on granular media. *Science*, 339(6126):1408–1412, 2013.

B. Liang and M. Sun. Nonlinear flight dynamics and stability of hovering model insects. *J R Soc Interface*, 10(85), Aug. 2013.

D. P. Looze, J. S. Freudenberg, J. H. Braslavsky, and R. H. Middleton. Tradeoffs and limitations in feedback systems. In W. S. Levine, editor, *The Control Systems Handbook*. CRC Press, Dec 2010.

H. Luo, R. Mittal, X. Zheng, S. A. Bielamowicz, R. J. Walsh, and J. K. Hahn. An immersed-boundary method for flow–structure interaction in biological systems with application to phonation. *J Comput Phys*, 227(22):9303–9332, 2008.

M. A. MacIver, N. M. Sharabash, and M. E. Nelson. Prey-capture behavior in gymnotid electric fish: Motion analysis and effects of water conductivity. *J Exp Biol*, 204(3):543–557, 2001.

M. S. Madhav, S. A. Stamper, E. S. Fortune, and N. J. Cowan. Closed-loop stabilization of the jamming avoidance response reveals its locally unstable and globally nonlinear dynamics. *J Exp Biol*, 216(22):4272–4284, 2013.

J. Matsubara and W. Heiligenberg. How well do electric fish electrolocate under jamming. *J Comp Physiol*, 125:285–290, 1978.

J. C. Maxwell. On governors. *Proc Royal Soc London*, pages 270–283, 1867.

W. Metzner. The jamming avoidance response in Eigenmannia is controlled by two separate motor pathways. *J Neurosci*, 13(5): 1862–78, May 1993.

R. H. Middleton. Trade-offs in linear control system design. *Automatica*, 27(2):281–292, 1991.

L. A. Miller, D. I. Goldman, T. L. Hedrick, E. D. Tytell, Z. J. Wang, J. Yen, and S. Alben. Using computational and mechanical models to study animal locomotion. *Integr Comp Biol*, 52 (5):553–575, 2012.

R. Mittal. Computational modeling in biohydrodynamics: trends, challenges, and recent advances. *Oceanic Engineering, IEEE Journal of*, 29(3):595–604, 2004.

J.-M. Mongeau, A. Demir, J. Lee, N. J. Cowan, and R. J. Full. Locomotion- and mechanics-mediated tactile sensing: antenna reconfiguration simplifies control during high-speed navigation in cockroaches. *J Exp Biol*, 216(24):4530–4541, 2013.

M. A. Nicolelis. *Methods for neural ensemble recordings*. CRC press, second edition, 2008.

H. F. Nijhout and M. C. Reed. Homeostasis and dynamic stability of the phenotype link robustness and plasticity. *Integr Comp Biol*, 2014. In review.

K. Nishikawa, A. A. Biewener, P. Aerts, A. N. Ahn, H. J. Chiel, M. A. Daley, T. L. Daniel, R. J. Full, M. E. Hale, T. L. Hedrick, A. K. Lappin, T. R. Nichols, R. D. Quinn, R. A. Satterlie, and B. Szymik. Neuromechanics: an integrative approach for understanding motor control. *Integr Comp Biol*, 47(1):16–54, Jul 2007.

D. C. O'Carroll, N. J. Bidweii, S. B. Laughlin, and E. J. Warrant. Insect motion detectors matched to visual ecology. *Nature*, 382 (6586):63–66, July 1996.

J. W. S. Pringle. The gyroscopic mechanism of the halteres of diptera. *Philos Trans R Soc B*, 233(602):347–384, Nov. 1948.

S. Revzen and J. M. Guckenheimer. Finding the dimension of slow dynamics in a rhythmic system. *J R Soc Interface*, 2011. doi: 10.1098/rsif.2011.0431. URL `http://rsif.royalsocietypublishing.org/content/early/2011/09/20/rsif.2011.0431.abstract`.

C. T. Richards. Building a robotic link between muscle dynamics and hydrodynamics. *J Exp Biol*, 214(14):2381–2389, June 2011.

C. T. Richards and C. J. Clemente. A bio-robotic platform for integrating internal and external mechanics during muscle-powered swimming. *Bioinspir Biomim*, 7(1):016010, Feb. 2012.

L. Ristroph, G. Ristroph, S. Morozova, A. J. Bergou, S. Chang, J. Guckenheimer, Z. J. Wang, and I. Cohen. Active and passive stabilization of body pitch in insect flight. *J R Soc Interface*, 10 (85):20130237, 2013.

G. J. Rose and J. G. Canfield. Discrimination of the sign of frequency differences by *Sternopygus*, an electric fish without a jamming avoidance response. *J Comp Physiol A*, 168:461–467, 1991.

G. J. Rose and J. G. Canfield. Longitudinal tracking responses of *Eigenmannia* and *Sternopygus*. *J Comp Physiol A*, 173:698–700, 1993a.

G. J. Rose and J. G. Canfield. Longitudinal tracking responses of the weakly electric fish, *Sternopygus*. *J Comp Physiol A*, 171 (6):791–798, Jan. 1993b.

E. Roth, K. Zhuang, S. A. Stamper, E. S. Fortune, and N. J. Cowan. Stimulus predictability mediates a switch in locomotor smooth pursuit performance for *Eigenmannia virescens*. *J Exp Biol*, 214(7):1170–1180, Apr. 2011.

E. Roth, S. Sponberg, and N. J. Cowan. A comparative approach to closed-loop computation. *Curr Opin Neurobiol*, 25:54–62, April 2014.

R. Ruiz-Torres, O. M. Curet, G. V. Lauder, and M. A. MacIver. Kinematics of the ribbon fin in hovering and swimming of the electric ghost knifefish. *J Exp Biol*, 216(5):823–834, 2013.

S. P. Sane, A. Dieudonne, M. A. Willis, and T. L. Daniel. Antennal mechanosensors mediate flight control in moths. *Science*, 315 (5813):863–866, Feb. 2007.

R. Schafer and T. V. Sanchez. Antennal sensory system of the cockroach, Periplaneta americana: postembryonic development and morphology of the sense organs. *J Comp Neurol*, 149:335–354, 1973.

D. Schaller. Antennal sensory system of Periplaneta americana L. *Cell and tissue research*, 191(1):121–139, 1978.

H. Scheich, T. H. Bullock, and R. H. Hamstra. Coding properties of two classes of afferent nerve fibers: high-frequency electroreceptors in the electric fish, eigenmannia. *J Neurophysiol*, 36(1): 39–60, 1973.

J. Schmitt and P. Holmes. Mechanical models for insect locomotion: dynamics and stability in the horizontal plane I. Theory. *Biol Cybern*, 83:501–515, 2000.

S. Sefati, E. S. Fortune, and N. J. Cowan. Counter-propogating waves in the ribbon fin of *Eigenmannia virescens* enhance maneuverability. In *Soc Int Comp Biol*, 2010.

S. Sefati, I. Neveln, M. A. MacIver, E. S. Fortune, and N. J. Cowan. Counter-propagating waves enhance maneuverability and stability: a bio-inspired strategy for robotic ribbon-fin propulsion. In *Proc IEEE Int Conf Biomed Robot Biomech.*, pages 1620–1625. IEEE, 2012.

S. Sefati, I. D. Neveln, E. Roth, T. Mitchell, J. B. Snyder, M. A. MacIver, E. S. Fortune, and N. J. Cowan. Mutually opposing forces during locomotion can eliminate the tradeoff between maneuverability and stability. *Proc Nat Acad Sci*, 110(47):18798–18803, 2013.

A. A. Shirgaonkar, O. M. Curet, N. A. Patankar, and M. A. Maciver. The hydrodynamics of ribbon-fin propulsion during impulsive motion. *J Exp Biol*, 211(Pt 21):3490–3503, Nov. 2008.

S. Sponberg and T. L. Daniel. Abdicating power for control: a precision timing strategy to modulate function of flight power muscles. *Proc Royal Soc B*, 279(1744):3958–3966, Oct. 2012.

S. Sponberg and R. J. Full. Neuromechanical response of musculoskeletal structures in cockroaches during rapid running on rough terrain. *J Exp Biol*, 211(3):433–446, Feb. 2008.

S. Sponberg, T. Libby, C. H. Mullens, and R. J. Full. Shifts in a single muscle's control potential of body dynamics are determined by mechanical feedback. *Philos Trans R Soc B*, 366(1570): 1606–1620, May 2011a.

S. Sponberg, A. J. Spence, C. H. Mullens, and R. J. Full. A single muscle's multifunctional control potential of body dynamics for postural control and running. *Philos Trans R Soc B*, 366(1570): 1592–1605, May 2011b.

S. Sponberg, A. J. Spence, C. H. Mullens, and R. J. Full. A single muscle's multifunctional control potential of body dynamics for postural control and running. *Philos Trans R Soc B*, 366(1570): 1592–1605, May 2011c.

S. A. Stamper, J. A. Simmons, C. M. Delong, and R. Bragg. Detection of targets colocalized in clutter by big brown bats (Eptesicus fuscus). *J Acoust Soc Am*, 124(1):667–673, Jul 2008.

S. A. Stamper, E. Carrera-G, E. W. Tan, V. Fugère, R. Krahe, and E. S. Fortune. Species differences in group size and electrosensory interference in weakly electric fishes: implications for electrosensory processing. *Behav Brain Res*, 207(2):368–376, 2010.

S. A. Stamper, M. S. Madhav, N. J. Cowan, and E. S. Fortune. Beyond the jamming avoidance response: Weakly electric fish respond to the envelope of social electrosensory signals. *J Exp Biol*, 215(23):4196–4207, 2012a.

S. A. Stamper, E. Roth, N. J. Cowan, and E. S. Fortune. Active sensing via movement shapes spatiotemporal patterns of sensory feedback. *J Exp Biol*, 215(9):1567–1574, 2012b.

M. Szwed, K. Bagdasarian, and E. Ahissar. Encoding of vibrissal active touch. *Neuron*, 40(3):621–630, 2003.

G. K. Taylor and H. G. Krapp. Sensory systems and flight stability: What do insects measure and why? In *Insect Mechanics and Control*, volume Volume 34, pages 231–316. Academic Press, 2007.

J. C. Theobald, E. J. Warrant, and D. C. O'Carroll. Wide-field motion tuning in nocturnal hawkmoths. *Proc Royal Soc B*, 277 (1683):853–860, Mar. 2010.

S. J. Thomas, R. R. Harrison, A. Leonardo, and M. S. Reynolds. A battery-free multichannel digital neural/emg telemetry system for flying insects. *Biomedical Circuits and Systems, IEEE Transactions on*, 6(5):424–436, 2012.

E. D. Tytell, C.-Y. Hsu, T. L. Williams, A. H. Cohen, and L. J. Fauci. Interactions between internal forces, body stiffness, and fluid environment in a neuromechanical model of lamprey swimming. *Proc Nat Acad Sci*, 107(46):19832–19837, Nov. 2010.

E. D. Tytell, P. Holmes, and A. H. Cohen. Spikes alone do not behavior make: why neuroscience needs biomechanics. *Curr Opin Neurobiol*, 21(5):816–822, Oct. 2011.

A. R. Wallace. On the tendency of varieties to depart indefinitely from the original type. *Proc Royal Soc*, pages 270–283, 1858.

Z. J. Wang, J. M. Birch, and M. H. Dickinson. Unsteady forces and flows in low reynolds number hovering flight: two-dimensional computations vs robotic wing experiments. *J Exp Biol*, 207(3): 449–460, 2004.

E. Warrant and M. Dacke. Vision and visual navigation in nocturnal insects. *Annu Rev Entomol*, 56:239–254, 2011.

N. Wiener. *Cybernetics: or Control and Communication in the Animal and the Machine*. John Wiley and Sons, Inc., New York, 1948.

J. H. Wu and M. Sun. Floquet stability analysis of the longitudinal dynamics of two hovering model insects. *J R Soc Interface*, 9 (74), Sept. 2012.

F. Zhang, L.-P. Wang, M. Brauner, J. F. Liewald, K. Kay, N. Watzke, P. G. Wood, E. Bamberg, G. Nagel, A. Gottschalk, and K. Deisseroth. Multimodal fast optical interrogation of neural circuitry. *Nature*, 446(7136):633–639, Apr 2007.